

How a brane cosmological constant can trick us into thinking that $w < -1$

Arthur Lue* and Glenn D. Starkman†

*Center for Education and Research in Cosmology and Astrophysics
Department of Physics, Case Western Reserve University, Cleveland, OH 44106-7079
and CERN Physics Department, Theory Division, CH-1211 Geneva 23, Switzerland*

Abstract

Observations exploring the contemporary cosmic acceleration have sparked interest in dark energy models possessing equations of state with $w < -1$. We review how the cosmic expansion history of a Dvali–Gabadadze–Porrati (DGP) braneworld model with a standard brane cosmological constant can mimic that of ordinary 4-dimensional gravity with $w < -1$ “phantom” dark energy for observationally relevant redshifts. We reinterpret the effective phantom nature of the dark energy as arising from dynamical-screening of the brane cosmological constant in DGP. This unusual variety of expansion history is thus possible without violating the null-energy condition, without ghosts and without any big rip, in a model which seems no more contrived than most evolving dark energy models. We indicate ways by which one may observationally test this effective $w < -1$ possibility, and differentiate it from “ordinary” phantom dark-energy.

*E-mail: lue@cern.ch

†E-mail: glenn.starkman@cern.ch

I. INTRODUCTION

The discovery of a contemporary cosmic acceleration [1,2] is one of the most profound scientific observations of the 20th century. The conventional explanation is some unknown smooth dark energy component that possesses a sufficiently negative pressure. Then, the Friedmann equation for a spatially flat Universe becomes

$$H^2 = \frac{8\pi G}{3} (\rho_M + \rho_{DE}) \ , \quad (1.1)$$

the dark energy has an equation of state w , where

$$\rho_{DE}(t) = \rho_{DE}^0 a^{-3(1+w)} \ . \quad (1.2)$$

The quantity $a(t)$ is the scale factor of the Universe. Dark energy composed of just a cosmological constant ($w = -1$) is fully consistent with existing observational data. However, that same data also suggests that if one is guided solely by the expansion history of the universe, $w < -1$ remains a viable possibility, and is slightly favored [3] (though with little statistical significance).

Regardless of the strength of this inference, $w < -1$ remains a logical and intriguing possibility. Many attempts have been made to find dark energy models that allow so-called phantom dark energy: dark energy with $w < -1$ [4–11]. Recently, Carroll, De Felice and Trodden [12] articulated the possibility that Friedmann equation Eq. (1.1) does not describe nature and that one may obtain a $w_{\text{eff}} < -1$ through a modification of the Friedmann equation rather than through an unusual (and often pathological) form of dark energy. They restricted their approach to Brans–Dicke–type theories and found “that only highly contrived models would lead observers to measure $w < -1$.”

Nevertheless, it is already known how one may achieve this goal within the context of modified-gravity theories. Several years ago, Sahni and Shtanov pointed out how to reproduce $w_{\text{eff}} < -1$ cosmologies [13] in the well-studied Dvali, Gabadadze and Porrati (DGP) braneworld theory [14–16] using the governing cosmological equations first developed by Deffayet [17]. This braneworld model is a complete theory, contains only matter satisfying the null-energy condition, exhibits no ghosts in the classical field theory, avoids a future big rip and requires no contrived system parameters.¹ In this note, we review the features of this braneworld model, taking a somewhat different approach for instructive purposes. We then discuss signatures by which one may identify this $w_{\text{eff}} < -1$ possibility observationally and distinguish it from “ordinary” phantom dark energy.

¹There is currently a debate regarding *quantum* strong coupling in this theory [18]. While this subject remains an important issue to resolve, it has no bearing on the present context where we are solely concerned with classical field evolution.

II. SCREENING OF THE COSMOLOGICAL CONSTANT

The DGP model is a 5-dimensional braneworld with modified gravity and thus a modified Friedmann equation [17]. Until now, phenomenological emphasis has been put on the possibility in this model of “self-acceleration” – acceleration with *no* negative-pressure energy-momentum component. Observationally, this leads to $w_{\text{eff}} > -1$, and this self-accelerating model has been studied extensively [17, 19–24]. However, there is another cosmological phase besides the self-acceleration one [17]. If one allows the most trivial negative-pressure energy-momentum component (a cosmological constant) then one can have $w_{\text{eff}} < -1$ [13]. We will rely on well-known governing equations already set out in the DGP literature, and will liberally appeal to that literature for details.

We invoke the following scenario: ordinary particles and fields, other than the graviton, are confined to a three-dimensional brane hypersurface embedded in a five-dimensional, infinite-volume Minkowski bulk [14]. The bulk is empty; all energy-momentum is isolated on the brane. The graviton is metastable on the brane. The action is [14]:

$$S_{(5)} = -\frac{1}{16\pi}M^3 \int d^5x \sqrt{-g} R + \int d^4x \sqrt{-g^{(4)}} \mathcal{L}_m . \quad (2.1)$$

M is the fundamental five-dimensional Planck scale. The first term in $S_{(5)}$ is the Einstein-Hilbert action in five dimensions for a five-dimensional metric g_{AB} (bulk metric) with Ricci scalar R and determinant g . The metric $g_{\mu\nu}^{(4)}$ is the induced (four-dimensional) metric on the brane, and $g^{(4)}$ is its determinant. An intrinsic curvature term is added to the brane action [14]:

$$-\frac{1}{16\pi}M_P^2 \int d^4x \sqrt{-g^{(4)}} R^{(4)} . \quad (2.2)$$

Here, M_P is the observed four-dimensional Planck scale (see [14–16] for details). Equation (2.2) is the Einstein-Hilbert action for the induced brane metric $g_{\mu\nu}^{(4)}$, $R^{(4)}$ being its scalar curvature.

Gravity in this model appears four-dimensional Einstein at short distances, while at larger distances, gravity evaporates into the bulk and is able to “probe” the extra dimension. The scale that governs that crossover is

$$r_0 = \frac{M_P^2}{2M^3} . \quad (2.3)$$

The cosmology on the brane for a spatially-flat brane Universe follows Deffayet’s modified Friedmann equation [17]

$$H^2 \pm \frac{H}{r_0} = \frac{8\pi G}{3} \rho(t) , \quad (2.4)$$

where $H(t) = \dot{a}/a$ is the Hubble parameter. The quantity $\rho(t)$ is the density of all energy momentum on the brane. The two choices of sign represent two distinct cosmological phases. The celebrated self-accelerating phase [17,19–21] corresponds to the lower sign, while we refer to the upper sign as the Minkowski cosmological phase. A natural geometric interpretation exists for both phases [17,25], but here we focus on the Minkowski phase. The key to this phase is that the brane is extrinsically curved in such a way that shortcuts through the bulk allow gravity to screen the effects of the brane energy-momentum contents at Hubble parameters $H \sim r_0^{-1}$. This is not the case for the self-accelerating phase.

Take the observed Universe (which is our brane) to have a homogeneous distribution of pressureless matter as well as a cosmological constant. Then,

$$H^2 + \frac{H}{r_0} = \frac{8\pi G}{3}\rho_M(t) + \lambda. \quad (2.5)$$

One may infer an effective dark energy component, c.f. Eq. (1.1), of the form

$$\frac{8\pi G}{3}\rho_{DE}^{\text{eff}} = \lambda - \frac{H}{r_0}. \quad (2.6)$$

Since $H(t)$ is a decreasing function of time, this effective dark energy component is increasing with time, the hallmark of a $w_{\text{eff}} < -1$ component. Indeed, one can easily define a w_{eff} using Eq. (1.1) and see that during the epoch when dark energy dominates, $w_{\text{eff}} < -1$. The second term in Eq. (2.6) is the effect of gravitational screening of the cosmological constant. As the universe evolves, that screening becomes increasingly ineffective and the effective dark-energy density appears to *increase*. The degree to which this effective dark-energy component grows with time is modulated by r_0 and in the limit $r_0 \rightarrow \infty$, we recover the standard Λ CDM cosmology. For any value of r_0 , $w_{\text{eff}} \rightarrow -1$ with time and a pure cosmological-constant Universe is restored in the distant future.²

III. EXPANSION HISTORY

Equations (2.5) and (2.6) imply that the cosmic expansion history does not strictly follow that for a conventional constant- w dark energy model. The precise time evolution of the Hubble parameter for this screened-cosmological-constant DGP may be determined explicitly from Eq. (2.5). Rewriting $H(t)$ in terms of more phenomenological quantities, we have

²In the past, when ρ_M dominates, the definition for w_{eff} breaks down, particularly when $\rho_{DE}^{\text{eff}} = 0$. This anomaly is an artifact of the definition: the evolution of $H(t)$ remains well-behaved throughout time.




FIG. 1. $d_L^w(z)/d_L^{\Lambda\text{CDM}}(z)$ and $d_L^{\lambda\text{DGP}}(z)/d_L^{\Lambda\text{CDM}}(z)$ for a variety of models. The reference model is the best-fit flat ΛCDM , with $\Omega_M^0 = 0.27$. The dotted curves are for the constant- w models with $(\Omega_M^w, w) = (0.28, -1.02), (0.294, -1.05)$, and $(0.316, -1.1)$ from top to bottom. The quantity Ω_M^w is chosen to match the ΛCDM model at low- z for a given w . The solid curves are for the screened-cosmological-constant DGP models with $(\Omega_M^{\lambda\text{DGP}}, r_0 H_0) = (0.292, 5.9), (0.34, 1.83)$, and $(0.42, 0.86)$ from top to bottom where the parameters were chosen to best mimic the corresponding w -models.

$$\frac{H(z)}{H_0} = \frac{1}{2} \left[-\frac{1}{r_0 H_0} + \sqrt{\left(2 + \frac{1}{r_0 H_0}\right)^2 + 4\Omega_M^0((1+z)^3 - 1)} \right], \quad (3.1)$$

where the redshift, z , today's Hubble parameter, H_0 , and $\Omega_M^0 = 8\pi G\rho_M^0/3H_0^2$ are all defined in the usual way. Given H_0 , there are two dimensionless parameters that may be adjusted phenomenologically: Ω_M^0 and $r_0 H_0$. This is similar to the conventional dark-energy picture, Eq. (1.1), which also has two phenomenological parameters: Ω_M^0 and w .

When comparing the expansion histories $a(t)$ of specific examples of screened-cosmological-constant DGP models (λDGP) with examples of constant- w dark-energy models, we refer to the Ω_M -parameter of the first as $\Omega_M^{\lambda\text{DGP}}$ and the second as Ω_M^w . For models with closely corresponding histories, these will not be the same. The luminosity-distance takes the standard form in spatially flat cosmologies:

$$d_L^{\lambda\text{DGP}}(z) = (1+z) \int_0^z \frac{dz}{H(z)}, \quad (3.2)$$

using Eq. (3.1). In Fig. 1, we compare this distance with the luminosity distance for a constant- w dark-energy model

$$d_L^w(z) = (1+z) \int_0^z \frac{H_0^{-1} dz}{\sqrt{\Omega_M^w(1+z)^3 + (1-\Omega_M^w)(1+z)^{3(1+w)}}}, \quad (3.3)$$

both normalized to the best-fit ΛCDM model. Figure 1 demonstrates that while the expansion histories of the currently considered model and constant- w dark-energy models are not identical, they are for all practical purposes indistinguishable using luminosity-distance

data. One should not put too much weight onto the particular values of $\Omega_M^{\lambda DGP}$ in Fig. 1 as we should be more interested in having this λ DGP model best match the data, rather than some w dark-energy cosmology. For a more thorough analysis of best-fit parameters of this model, see Alam and Sahni [26].

IV. OBSERVATIONAL SIGNATURES

Up until now, we have reviewed what exists in the literature (though presented in a slightly different way) but, for whatever reason, remains largely unappreciated by the cosmology community. However, we can now ask important questions about observational constraints of this λ DGP model, and in particular how one may differentiate this modified-gravity model from dark-energy models that may mimic the same cosmic expansion history.

The major tests of this theory come from constraints on the modification of the gravitational force law. Because this model is in the class of more general DGP models, it does not suffer from the same constraints as found for Brans–Dicke theories of gravity. Extra scalar degrees of freedom are suppressed at distance scales much shorter than $r_0 \sim H_0^{-1}$, thus we expect current observational constraints on the form of gravity are not spoiled (see the discussion in [27–30,22,23]).

Generalizing the analysis found in Refs. [22,24] we may ascertain the modification of the Schwarzschild metric for a spherically symmetric source for any background cosmology. The metric on the brane

$$ds^2 = g_{00}dt^2 - g_{rr}dr^2 - r^2d\Omega . \quad (4.1)$$

For a cosmological background with *arbitrary* evolution $H(t)$, we find that

$$rg'_{00}(t,r)|_{\text{brane}} = \frac{R_g}{r} [1 + \Delta(r)] - 2(H^2 + \dot{H})r^2 \quad (4.2)$$

$$g_{rr}(t,r)|_{\text{brane}} = 1 + \frac{R_g}{r} [1 - \Delta(r)] + H^2r^2 , \quad (4.3)$$

where dot now denotes differentiation with respect to t and prime denotes differentiation with respect to r . Note that the cosmological background contribution is included in these metric components. The quantity R_g is defined as

$$R_g(r) = 8\pi G \int_0^r dr r^2 \delta\rho(r) , \quad (4.4)$$

where $\delta\rho(r)$ is density of matter in excess of the background cosmological density; $\Delta(r)$ is defined as

$$\Delta(r) = \frac{3\beta r^3}{4r_0^2 R_g} \left[\sqrt{1 + \frac{8r_0^2 R_g}{9\beta^2 r^3}} - 1 \right] ; \quad (4.5)$$

and

$$\beta = 1 \pm 2r_0 H \left(1 + \frac{\dot{H}}{3H^2} \right) . \quad (4.6)$$

Just as for the modified Friedmann equation, Eq. (2.4), there is a sign degeneracy in Eq. (4.6). The lower sign corresponds to the self-accelerating cosmologies. For the cosmological phase in which we are interested (λ DGP), we take the upper (positive) sign.³

A. Solar system

For solar system measurements, one may take a limit in Eqs. (4.2) and (4.3) where $r^3 \ll r_0^2 R_g$. Then, there are only small corrections to the usual Schwarzschild metric. To leading-order, the immediate consequence of this modification of the Schwarzschild metric is that all orbiting bodies suffer a universal anomalous precession of [22]

$$\frac{d}{dt} \Delta\phi = -\frac{3}{8r_0} . \quad (4.7)$$

If $r_0 H_0 = 1$, then the precession rate is about $3 \mu\text{as}/\text{year}$. Note that this rate is independent of the source mass and only depends on the distance scale r_0 which characterizes the scale at which bulk effect manifest themselves in the Universe. Current solar system constraints are around $10 \mu\text{as}/\text{year}$, and improvements can be expected over the next decade [22,23]. It is intriguing that one may gain insight into physics on the largest of cosmological scales from local tests of gravity within our solar system.

B. Large scale structure

One may now also invoke tests from large scale structure. For linear perturbations, Eqs. (4.2) and (4.3) yield a metric for a linearized scalar-tensor theory with Brans–Dicke parameter

$$\omega = \frac{3}{2}(\beta - 1) . \quad (4.8)$$

Note this is only true for weak, linear perturbations around the cosmological background, and that for solar-system tests one must refer to the previous subsection. However, for the linear growth of structure, this is perfectly adequate. Nonrelativistic matter perturbations with $\delta(t) = \delta\rho/\rho$ evolve via

³There may be evidence of alternative nonperturbative phases of compact matter sources [31].

$$\ddot{\delta} + 2H\dot{\delta} = 4\pi G\rho \left(1 + \frac{1}{3\beta}\right) \delta , \quad (4.9)$$

implying the effect of λ DGP on linear matter perturbations is that self-attraction of overdensities is governed by an evolving G_{eff} :

$$G_{\text{eff}} = G \left[1 + \frac{1}{3\beta}\right] . \quad (4.10)$$

This effective Newton's constant runs from G at early times to values larger than G as redshift $z \rightarrow 0$. This mild strengthening of the self-attraction of density perturbations will increase perturbation growth, even beyond that one would expect from that from a dark energy model that mimics the same cosmic expansion history.

For example, by numerically evolving Eq. (4.9), one finds for the corresponding parameters $(\Omega_M^{\lambda DGP}, r_0 H_0) = (0.3, 3.77), (0.35, 1.41)$ and $(0.4, 0.870)$, we get anomalously high growth of λ DGP (beyond that one would get for a dark-energy model with the same cosmic expansion history) of 1.5%, 4.6% and 9.3%, respectively. This discrepancy of linear growth is roughly proportional to $1 - w_{\text{eff}}$ and provides an intriguing handle by which to differentiate between this modified-gravity scenario and corresponding dark-energy ones.

V. CONCLUDING REMARKS

We illustrate in this note how one may mimic a $w < -1$ dark-energy cosmology using just a brane cosmological constant in DGP in the Minkowski cosmological phase. The effect of this braneworld scenario is that leakage of gravity off the brane at late times screens the effect of the cosmological constant. This screening attenuates with time, yielding a dark-energy density that appears to grow as the universe evolves, thus providing a cosmic evolution with $w_{\text{eff}} < -1$. There are no currently catastrophic observational constraints on this model, though data in the near future may provide some intriguing tests. There may be an important roll for modified gravity to play in understanding the details of what drives today's cosmic acceleration.

ACKNOWLEDGMENTS

The authors thank G. Gabadadze and R. Scoccimarro for helpful communications. We thank the CERN Theory Division for its hospitality. This work is sponsored by DOE Grant DEFG0295ER40898, the CWRU Office of the Provost, and CERN. GDS thanks Maplesoft for the use of Maple V.

REFERENCES

- [1] S. Perlmutter *et al.* [Supernova Cosmology Project Collaboration], *Astrophys. J.* **517**, 565 (1999).
- [2] A. G. Riess *et al.* [Supernova Search Team Collaboration], *Astron. J.* **116**, 1009 (1998).
- [3] A. G. Riess *et al.* [Supernova Search Team Collaboration], *Astrophys. J.* **607**, 665 (2004).
- [4] R. R. Caldwell, *Phys. Lett. B* **545**, 23 (2002).
- [5] T. Barreiro, E. J. Copeland and N. J. Nunes, *Phys. Rev. D* **61**, 127301 (2000).
- [6] T. Chiba, T. Okabe and M. Yamaguchi, *Phys. Rev. D* **62**, 023511 (2000).
- [7] V. K. Onemli and R. P. Woodard, *Class. Quant. Grav.* **19**, 4607 (2002).
- [8] S. M. Carroll, M. Hoffman and M. Trodden, *Phys. Rev. D* **68**, 023509 (2003).
- [9] A. Melchiorri, L. Mersini, C. J. Odman and M. Trodden, *Phys. Rev. D* **68**, 043509 (2003).
- [10] J. M. Cline, S. y. Jeon and G. D. Moore, arXiv:hep-ph/0311312.
- [11] V. K. Onemli and R. P. Woodard, arXiv:gr-qc/0406098.
- [12] S. M. Carroll, A. De Felice and M. Trodden, arXiv:astro-ph/0408081.
- [13] V. Sahni and Y. Shtanov, *JCAP* **0311**, 014 (2003).
- [14] G. Dvali, G. Gabadadze and M. Porrati, *Phys. Lett. B* **485**, 208 (2000).
- [15] G. R. Dvali, G. Gabadadze, M. Kolanovic and F. Nitti, *Phys. Rev. D* **64**, 084004 (2001).
- [16] G. R. Dvali, G. Gabadadze, M. Kolanovic and F. Nitti, *Phys. Rev. D* **65**, 024031 (2002).
- [17] C. Deffayet, *Phys. Lett. B* **502**, 199 (2001).
- [18] M. A. Luty, M. Porrati and R. Rattazzi, *JHEP* **0309**, 029 (2003); V. A. Rubakov, arXiv:hep-th/0303125; A. Nicolis and R. Rattazzi, *JHEP* **0406**, 059 (2004); G. Gabadadze and M. Shifman, *Phys. Rev. D* **69**, 124032 (2004).
- [19] C. Deffayet, G. R. Dvali and G. Gabadadze, *Phys. Rev. D* **65**, 044023 (2002).
- [20] C. Deffayet, S. J. Landau, J. Raux, M. Zaldarriaga and P. Astier, *Phys. Rev. D* **66**, 024019 (2002).
- [21] J. S. Alcaniz, *Phys. Rev. D* **65**, 123514 (2002); D. Jain, A. Dev and J. S. Alcaniz, *Phys. Rev. D* **66**, 083511 (2002); J. S. Alcaniz, D. Jain and A. Dev, *Phys. Rev. D* **66**, 067301 (2002).
- [22] A. Lue and G. Starkman, *Phys. Rev. D* **67**, 064002 (2003).
- [23] G. Dvali, A. Gruzinov and M. Zaldarriaga, *Phys. Rev. D* **68**, 024012 (2003).
- [24] A. Lue, R. Scoccimarro and G. D. Starkman, *Phys. Rev. D* **69**, 124015 (2004).
- [25] A. Lue, *Phys. Rev. D* **67**, 064004 (2003).
- [26] U. Alam and V. Sahni, arXiv:astro-ph/0209443.
- [27] C. Deffayet, G. R. Dvali, G. Gabadadze and A. I. Vainshtein, *Phys. Rev. D* **65**, 044026 (2002).
- [28] A. Lue, *Phys. Rev. D* **66**, 043509 (2002).
- [29] A. Gruzinov, arXiv:astro-ph/0112246.
- [30] M. Porrati, *Phys. Lett. B* **534**, 209 (2002).
- [31] G. Gabadadze and A. Iglesias, arXiv:hep-th/0407049.